

## SEQUENCE LISTING

<110> KISIELOW, Malgorzata Anna  
KLEINER, Sandra  
NAGAMINE, Yoshikuni

<120> METHODS OF OBTAINING ISOFORM SPECIFIC  
EXPRESSION IN MAMMALIAN CELLS

<130> 1-32330A/FMI

<150> GB 0201477.7

<151> 2002-01-23

<150> PCT/EP03/00611

<151> 2003-01-22

<160> 16

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> oligoribonucleotide pair used as siRNA; h/m-shc  
siRNA from nt 677-697 (in the PTB domain)

<400> 1

cuacuugguu cgguacauagg g

21

<210> 2

<211> 22

<212> RNA

<213> Artificial Sequence

<220>

<223> oligoribonucleotide pair used as siRNA; h/m-shc  
siRNA from nt 677-697 (in the PTB domain)

<400> 2

cauguaccga acccaaguag ga

22

<210> 3

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> oligonucleotide pair used for siRNA; h/m-shc  
siRNA, p66-shc siRNA (from nt 236-256 in the CH2  
domain)

<400> 3  
 gaaugagucu cugucaucgu c 21  
  
 <210> 4  
 <211> 21  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide pair used in siRNA; h/m-shc siRNA,  
 p66-shc siRNA (from nt 236-256 in the CH2 domain)  
  
 <400> 4  
 cgaugacaga gacucauucc g 21  
  
 <210> 5  
 <211> 30  
 <212> DNA  
 <213> Mus musculus  
  
 <220>  
 <221> primer\_bind  
 <222> (1)...(30)  
 <223> sense primer for mouse p46 ShcA cDNA  
  
 <400> 5  
 cggaattcat gggacctggg gtttcctact 30  
  
 <210> 6  
 <211> 30  
 <212> DNA  
 <213> Mus musculus  
  
 <220>  
 <221> primer\_bind  
 <222> (1)...(30)  
 <223> sense primer for mouse p52 ShcA cDNA  
  
 <400> 6  
 cggaattcat gaacaagctg agtggaggcg 30  
  
 <210> 7  
 <211> 36  
 <212> DNA  
 <213> Mus musculus  
  
 <220>  
 <221> primer\_bind  
 <222> (1)...(36)  
 <223> sense primer for mouse p66 ShcA cDNA  
  
 <400> 7  
 cggaattcat ggatcttcta ccccccaagc cgaagt 36  
  
 <210> 8  
 <211> 31  
 <212> DNA

<213> Mus musculus  
 <220>  
 <221> primer\_bind  
 <222> (1)...(31)  
 <223> common antisense primer for ShcA cDNAs  
 <400> 8  
 cggaattcac actttccgat ccacgggttg c 31  
 <210> 9  
 <211> 40  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> overlapping oligonucleotide pair for insertion  
 into the KpnI-EcoRI site of pcDNA3  
 <400> 9  
 cccaccatgg cttaccata cgatgttcca gattacgctg 40  
 <210> 10  
 <211> 48  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> overlapping oligonucleotide pair for insertion  
 into the KpnI-EcoRI site of pcDNA3  
 <400> 10  
 aattcagcga attctggaac atcgtatggg taagccatgg tggggtac 48  
 <210> 11  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> overlapping oligonucleotide pair used to mutate  
 methionine 65 (start site for p52) to leucine in  
 p66HA, resulting in p66HA-m1  
 <400> 11  
 ctcttcagg acctgaacaa gctgagt 27  
 <210> 12  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> overlapping oligonucleotide pair used to mutate  
 methionine 65 (start site for p52) to leucine in  
 p66HA, resulting in p66HA-m1

<400> 12	
cactcagctt gttcagggtcc tggaggag	28
<210> 13	
<211> 27	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to mutate the initiation sites for p46 in both p66HA-m1 and p52HA, resulting in p66HA-ML and p52HA-ML	
<400> 13	
ccaacgacaa agtcctggga cccgggg	27
<210> 14	
<211> 27	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to mutate the initiation sites for p46 in both p66HA-m1 and p52HA, resulting in p66HA-ML and p52HA-ML	
<400> 14	
ccccgggtcc caggactttg tcgttgg	27
<210> 15	
<211> 34	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to give p46HA-sm, p52HA-ML-sm and p66HA-ML-sm	
<400> 15	
ggggtttcct acttggtccg ctacatgggt tgtc	34
<210> 16	
<211> 34	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to give p46HA-sm, p52HA-ML-sm and p66HA-ML-sm	
<400> 16	
cacaacccat gtagcggacc aagtaggaaa cccc	34